

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Pro	vince, China
TELE: 86-510-8	FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>	
Email: info@dlde	velop.com.cn service@dldevelop.com.cn
Product name:	Anti-Mouse H-2 Monoclonal Antibody(PE/Cyanine7
	Conjugated)
Catalog:	DL22282F
cutility.	
Synonyms:	Mouse major histocompatibility complex (MHC)
Synonyms.	H-2, MHC I
	11-2, MITC 1
Daalamaanda	The M1/42 antihedry recets with the U.2 MUC class I
Background:	The M1/42 antibody reacts with the H-2 MHC class I
	alloantigens expressed on nucleated cells from mice
	of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC
	et al, 1981). MHC class I is involved in antigen
	presentation to T cells expressing CD3/TCR and
	CD8 proteins.
Form: Liquid	Isotype: Rat IgG2a, κ
Size: 25µg/100µg	Host: Rat



Reactivity: Mouse

Application: FCM

Concentration: 0.2 mg/mL

Conjugation: PE/Cyanine7

SwissProt:

Buffer:Phosphate buffered solution, pH 7.2, containing
0.09% stabilizer and 1% protein protectant.


Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu g/10^6$ cells in 100 μ L volume].

Data:

C57BL/6 murine splenocytes are stained with Anti-Mouse H-2 Monoclonal Antibody(PE/Cyanine7 Conjugated)[Used at 0.2 μ g/10⁶ cells dilution](filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.

DEVELOP



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.