

## Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District Wuxi Jiangsu Province, China TELE: 86-510-82732223 FAX: 86-510-82720101-8014 Web : www.dldevelop.com Email: info@dldevelop.com.cn service@dldevelop.com.cn Product name: Anti-Rat CD44H Monoclonal Antibody(PE Conjugated) **Catalog:** DL22288F **Synonyms:** Pgp-1, H-CAM, CD44s, CD44H **Background:** CD44 is an 80-95 kD cell surface glycoprotein. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. It is up-regulated when T cells and B cells are activated. It was reported that CD44 is a valuable marker for memory T cells. CD44 is an adhesion molecule involved in leukocyte adhesion and homing to lymphoid organs. The OX-49 antibody reacts with CD44H (known as CD44s) expressed on most leukocytes, except for a subset of B lymphocytes. The epitope recognized by OX-49 antibody has been

www.dldevelop.com



	the splice variant However it was 1 detect the CD44	on on both the standard, CD44s, and t, CD44v, isoforms of CD44. reported that OX-49 antibody cannot V isoform, possibly due to changes in the epitope.
Form: Liquid		<b>Isotype:</b> Mouse IgG2a, κ
Size:		Host: Mouse
50Tests/100Tests	s/100Tests×2	
Reactivity: Rat		Application: FCM
Concentration:	5 μL	Conjugation: PE
SwissProt:		

Buffer:Phosphate buffered solution, pH 7.2, containing<br/>0.09% stabilizer and 1% protein protectant.

#### **Recommended Use:**


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined

# **DEVELOP**.

for individual use.

### Data:

Rat splenocytes are stained with Anti-Rat CD44H Monoclonal Antibody(PE Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



### Storage:

Keep as concentrated solution.

Store at  $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.