

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Anti-Rat CD44H Monoclonal Antibody(APC

Conjugated)

Catalog: DL22289F

Synonyms: Pgp-1, H-CAM, CD44s, CD44H

Background: CD44 is an 80-95 kD cell surface glycoprotein. It is

> expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. It is

up-regulated when T cells and B cells are activated. It was reported that CD44 is a valuable marker for memory T cells. CD44 is an adhesion molecule involved in leukocyte adhesion and homing to lymphoid organs. The OX-49 antibody reacts with CD44H (known as CD44s) expressed on most leukocytes, except for a subset of B lymphocytes. The epitope recognized by OX-49 antibody has been



mapped to a region on both the standard, CD44s, and the splice variant, CD44v, isoforms of CD44. However it was reported that OX-49 antibody cannot detect the CD44V isoform, possibly due to conformational changes in the epitope.

Form: Liquid **Isotype:** Mouse IgG2a, κ

Size: Host: Mouse

50Tests/100Tests/100Tests×2

Reactivity: Rat **Application:** FCM

Concentration: 5 µL **Conjugation:** APC

SwissProt:

Phosphate buffered solution, pH 7.2, containing Buffer:

0.09% stabilizer and 1% protein protectant.

Recommended Use:


Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined



for individual use.

Data:

Rat splenocytes are stained with Anti-Rat CD44H Monoclonal Antibody(APC Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at 2~8°C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.