

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-801				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Anti-Rat CD45 Monoclonal Antibody(PE/Cyanine5			
	Conjugated)			
Catalog:	DL22306F			
Synonyms:	Receptor-type tyrosine-protein phosphatase C,Leukocyte common antigen, LCA, T200,			
	Ly-5,Ptprc			
Background:	CD45 is a 180-220 kD protein also known as			
leukocyte common antigen (LCA). It is a protein tyrosine phosphatase with multiple isoforms differing				
		as a result of alternative splicing of the extracellular		
	domain and glycosylation. CD45 is expressed on all			
hematopoietic cells except erythrocytes and platelets				
	isoform expression depends on cell type, activation			
state, and cell maturation. CD45 functions in signal				
transduction through T and B cell antigen receptors				



CD45 has been shown to interact with various		
proteins including galectin-1, CD2, CD3, and CD4.		
The OX-1 antibody has been shown to partially		
inhibit NK cell-mediated lysis of syngeneic tumor		
cells in vitro.		

Form: Liquid		Isotype: Mouse IgG1, κ
Size: 50Tests/100Tests.	/100Tests×2	Host: Mouse
Reactivity: Rat		Application: FCM
Concentration: 5 µL		Conjugation: PE/Cyanine5
SwissProt:	P04157	
Buffer:	Phosphate buffered solution, pH 7.2, containing	

Recommended Use:

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined


0.09% stabilizer and 1% protein protectant.

DEVELOP

for individual use.

Data:

Rat splenocytes are stained with Anti-Rat CD45 Monoclonal Antibody(PE/Cyanine5 Conjugated)(filled gray histogram). Unstained splenocytes (empty black histogram) are used as control.



Storage:

Keep as concentrated solution.

Store at $2\sim 8^{\circ}$ C and protected from prolonged exposure to light. Do not freeze.

This product is guaranteed up to one year from purchase.