

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

**Product name:** Rabbit anti-ATF4 Antibody

Catalog: DL90082A

**Synonyms:** CREB2; TXREB; Cyclic AMP-dependent

transcription factor ATF-4; cAMP-dependent

transcription factor ATF-4; Activating transcription factor 4; Cyclic AMP-responsive element-binding

protein 2; CREB-2; cAMP-responsive

element-binding protein 2; DNA-binding pr

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human ATF4.

The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit



Reactivity: Human, Mouse, Rat **Application:** WB, IHC

**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200)

**Entrez Gene:** 468/11911/79255

P18848/Q06507/Q9ES19 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

## **WB** description:

Western blot analysis of ATF4 expression in MCF7 (A), mouse brain (B), rat brain (C) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of ATF4 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.