

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name: Catalog:	Rabbit anti-Cası DL90147A	pase 6 p18 /	Antibody	
Synonyms:	MCH2; Caspase-6; CASP-6; Apoptotic protease Mch-2			
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Caspase 6 p18. The exact sequence is proprietary.			
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Mouse, Rat		Application: WB, IHC, IF/IC		
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000),		



IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	839/12368
SwissProt:	P55212/O08738/O35397
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Caspase 6 p18 expression in HeLa (A), mouse brain (B), rat kidney (C), NIH3T3 staurosporine-treated (D), Jurkat etoposide-treated (E) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of Caspase 6 p18 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Caspase 6 p18 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

