


| Manufacturer/Supplier: | | | | |
|---|---|---------------------------------------|----------------------|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | | |
| Wuxi Jiangsu Province, China | | | | |
| TELE: 86-510-82732223 | | FAX: | 86-510-82720101-8014 | |
| Web : <u>www.dldevelop.com</u> | | | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | | | |
| | | | | |
| Product name: | Rabbit anti-Cyclin F Antibody | | | |
| Catalog: | DL90172A | | • | |
| 8 | | | | |
| Synonyms: | FBX1; FBXO1; Cyclin-F; F-box only protein 1 | | | |
| Synonyms. | | Cyclin 1,1 | box only protein 1 | |
| Immunogen: | KLH-conjugated synthetic peptide encompassing a | | | |
| Ininunogen. | sequence within the C-term region of human Cyclin | | | |
| | | | | |
| F. The exact sequence is proprietary. | | | | |
| . . | | C | | |
| Form: Liquid | | Concentr | ation: 1mg/mL | |
| | | | | |
| Size: 100 ul/50 ul | | Host: Rabbit | | |
| | | | | |
| Reactivity: Human, Mouse, Rat | | Applicati | on: WB, IHC | |
| | | | | |
| Clonality: Polyclonal | | Dilution: WB (1/500 - 1/1000), | | |
| | | IHC (1/10 | 0 - 1/200) | |
| | | | | |

DEVELOP.

| Entrez Gene: | 899/12449/117524 |
|---------------|---|
| SwissProt: | P41002/P51944/Q8K4F8 |
| Purification: | The antibody was purified by immunogen affinity chromatography. |
| Buffer: | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |

WB description:

Western blot analysis of Cyclin F expression in HEK293T (A), mouse liver (B), rat liver (C) whole cell lysates.



IHC description:

Immunohistochemical analysis of Cyclin F staining in human



breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

