

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn) [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Rabbit anti-E Cadherin Antibody

**Catalog:** DL90199A

**Synonyms:** CDH1; CDHE; UVO; Cadherin-1; CAM 120/80;  
Epithelial cadherin; E-cadherin; Uvomorulin;  
CD324; CDH2; CDHN; NCAD; Cadherin-2;  
CDw325; Neural cadherin; N-cadherin; CD325;  
CDH3; CDHP; Cadherin-3; Placental cadherin;  
P-cadherin; CDH4; Cadherin-4; Retinal cadher

**Immunogen:** KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human E Cadherin. The exact sequence is proprietary.

**Form:** Liquid

**Concentration:** 1mg/mL

**Size:** 100 ul/50 ul

**Host:** Rabbit

**Reactivity:** Human,Rat,Bovine,Dog,Monkey, Pig,Rabbit,Zebrafish  
**Application:** WB, IHC, IF/IC, IP, FC

**Clonality:** Polyclonal  
**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100)

**Entrez Gene:** 1017


**SwissProt:** P24941

**Purification:** The antibody was purified by immunogen affinity chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

**WB description:**

Western blot analysis of CDK2 expression in Jurkat (A) whole cell lysates.




### IHC description:


Immunohistochemical analysis of CDK2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### IF/ICC description:

Immunofluorescent analysis of CDK2 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



**IP description:**



**FC description:**

**Storage:**



Store at -20°C. Avoid repeated freeze / thaw cycles.