

Manufacturer/Su	pplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name: Catalog:	Rabbit anti-CDF DL90205A	X2 Antibod	у	
Synonyms:	CDKN2; Cyclin-dependent kinase 2; Cell division protein kinase 2; p33 protein kinase			
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CDK2. The exact sequence is proprietary.			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
<b>Reactivity:</b> Human,Bovine,Ch Rabbit	icken,Dog,Pig,	Application	on: WB, IHC, IF/IC, IP	



**Clonality:** Polyclonal


**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 -1/500), IP (1/10 - 1/100)

Entrez Gene:	1434/110750
SwissProt:	P55060/Q9ERK4
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of XPO2 expression in HeLa (A), NIH3T3 (B), SP2/0 (C), PC12 (D) whole cell lysates.

# **DEVELOP**




### **IHC description:**

Immunohistochemical analysis of XPO2 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### **IF/ICC description:**

Immunofluorescent analysis of XPO2 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

## **DEVELOP**.



**IP description:** 

**Storage:** 

Store at -20°C. Avoid repeated freeze / thaw cycles.

