

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Deve deve to a service	D-11:44: C-11	(-11.		
Product name:	Rabbit anti-Collagen 6 alpha 2 Antibody			
Catalog:	DL90256A			
Supervise	Collogon alpha	2(VI) shain		
Synonyms:	Collagen alpha-2(VI) chain			
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
8	sequence within the center region of human			
	Collagen 6 alpha		e	
	proprietary.	a 2. The exe	iet sequence is	
	proprietary			
Form: Liquid		Concentra	ation: 1mg/mL	
1			0	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Monkey		Applicati	on: WB, IHC, IF/IC, IP	

Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000),



IHC (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100)


Entrez Gene:	1292
SwissProt:	P12110
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Collagen 6 alpha 2 expression in HepG2 (A), HT29 (B) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of Collagen 6 alpha 2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Collagen 6 alpha 2 staining in HT29 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the prim



DEVELOP.



IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

