

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Lamin A/C Antibody			
Catalog:	DL90691A			
Synonyms:	LMN1; Prelamin	n-A/C		
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within	the center	region of human Lamin	
	A/C. The exact s	sequence is	proprietary.	
T T ' ' 1		C (· 1 / T	
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Size. 100 ul/30 ul		Host. Rac		
Reactivity:		Applicati	on: WB, IHC, IF/IC	
Human,Mouse,Rat,Chicken,Pig				
Clonality: Polyclonal		Dilution:	WB (1/500 - 1/1000),	

www.dldevelop.com



IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	4000/16905/60374
SwissProt:	P02545/P48678/P48679
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Lamin A/C expression in HeLa colchicine-treated (A), mouse liver (B), rat kidney (C) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of Lamin A/C staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Lamin A/C staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

