

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-c-Myc (pT58) Antibody			
Catalog:	DL90776A			
~				
Synonyms:	BHLHE39; Myc proto-oncogene protein; Class E			
			in 39; bHLHe39;	
	Proto-oncogene	e c-Myc; Tra	inscription factor p64	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
g	sequence within the N-term region of human c-Myc.			
	The exact sequence is proprietary.			
	The chuct sequ		liotui y.	
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity:		Applicati	on: WB, IHC, IP	
Human,Mouse,Rat,Bovine,Chicke				

DEVELOP.

n,Dog,Pig,Sheep,Zebrafish


Clonality: Polyclonal	Dilution: WB (1/500 - 1/1000),
	IHC (1/100 - 1/200), IP (1/10 -
	1/100)

Entrez Gene:	4609/17869/24577
SwissProt:	P01106/P01108/P09416
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of c-Myc (pT58) expression in A431 (A), HeLa (B) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of c-Myc (pT58) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

