

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

Product name: Rabbit anti-NDUFA8 Antibody

Catalog: DL90788A

Synonyms: NADH dehydrogenase [ubiquinone] 1 alpha

subcomplex subunit 8; Complex I-19kD; CI-19kD; Complex I-PGIV; CI-PGIV; NADH-ubiquinone

oxidoreductase 19 kDa subunit

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human NDUFA8. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Application: WB, IHC, IP



Human, Mouse, Rat, Bovine, Monke

y

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IP (1/10 -

1/100)

Entrez Gene: 4702/68375

P51970/O9DCJ5 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of NDUFA8 expression in HepG2 (A), MCF7 (B) whole cell lysates.





IHC description:

Immunohistochemical analysis of NDUFA8 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.