

Manufacturer/Su	pplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Rabbit anti-PLC gamma 1 Antibody		
Catalog:	DL90897A		
Synonyms:	PLC1; 1-phosphatidylinositol 4,5-bisphosphate phosphodiesterase gamma-1; PLC-148; Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-gamma-1		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
	sequence within the center region of human PLC		
	gamma 1. The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
	U (D 11)		
Size: 100 ul/50 ul	Host: Rabbit		




Reactivity:		Application: WB, IHC	
Human, Mouse, Rat, Bovine, Monke			
У			
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)	
Entrez Gene:	5335/18803/25738		
SwissProt:	P19174/Q62077/P10686		
Purification:	The antibody was purified by immunogen affinity chromatography.		
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.		

WB description:

Western blot analysis of PLC gamma 1 expression in HEK293T (A), HeLa (B), SP2/0 (C), PC12 (D) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of PLC gamma 1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.