

Manufacturer/Supplier:WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTDAddress: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi DistrictWuxi Jiangsu Province, ChinaTELE: 86-510-82732223FAX: 86-510-82720101-8014Web: www.dldevelop.comEmail: info@dldevelop.com.cnservice@dldevelop.com.cn			
Product name:	Rabbit anti-ERK1/2 Antibody		
Catalog:	DL90946A		
Synonyms:	MAPK3; ERK1; PRKM3; Mitogen-activated protein kinase 3; MAP kinase 3; MAPK 3; ERT2; Extracellular signal-regulated kinase 1; ERK-1; Insulin-stimulated MAP2 kinase; MAP kinase isoform p44; p44-MAPK; Microtubule-associated protein 2 kinase; p44-ERK1; MAPK1;		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
	sequence within the center region of human		
	ERK1/2. The exact sequence is proprietary.		
Form: Liquid Size: 100 ul/50 ul	Concentration: 1mg/mL Host: Rabbit		




Reactivity: Human,Mouse,Rat, n,Zebrafish	Bovine,Chicke	Application: WB, IHC, IP
Clonality: Polyclo	nal	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)
Entrez Gene:	26417; 26413/50689; 116590	
SwissProt:	P27361; P28482	/Q63844; P63085/P21708; P63086
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	1	Potassium phosphate, 0.87% , pH 7.3, 30% glycerol, and 0.01%

WB description:

Western blot analysis of ERK1/2 expression in MCF7 (A), NIH3T3 (B), PC12 (C) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of ERK1/2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

