

Manufacturer/Supplier:WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTDAddress: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi DistrictWuxi Jiangsu Province, ChinaTELE: 86-510-82732223FAX: 86-510-82720101-8014Web : www.dldevelop.comEmail: info@dldevelop.com.cn			
Product name:	Rabbit anti-JNK3 Antibody		
Catalog:	DL90948A		
Synonyms:	JNK3; JNK3A; PRKM10; SAPK1B; Mitogen-activated protein kinase 10; MAP kinase 10; MAPK 10; MAP kinase p49 3F12; Stress-activated protein kinase 1b; SAPK1b; Stress-activated protein kinase JNK3; c-Jun N-terminal kinase 3		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human JNK3.		
0			
	The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
<b>Size:</b> 100 ul/50 ul	Host: Rabbit		




Reactivity: Human, Mouse, Rat		Application: WB, IF/IC, IP
Clonality: Polyclonal		<b>Dilution:</b> WB (1/500 - 1/1000), IF/IC (1/100 - 1/500), IP (1/10 - 1/100)
Entrez Gene:	5602/26414/25272	
SwissProt:	P53779/Q61831/P49187	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

## **WB description:**


Western blot analysis of JNK3 expression in NIH3T3 (A), SKNSH (B), HeLa (C) whole cell lysates.

## **DEVELOP**



## **IF/ICC description:**

Immunofluorescent analysis of JNK3 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



**IP description:** 



## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.

