

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-NF-kappaB p65 (pS468) Antibody

Catalog: DL90998A

Synonyms: NFKB3; Transcription factor p65; Nuclear factor

NF-kappa-B p65 subunit; Nuclear factor of kappa

light polypeptide gene enhancer in B-cells 3

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the C-term region of human

NF-kappaB p65. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Application: WB, IHC, IP

Human, Mouse, Rat, Bovine, Monke



y,Pig

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IP (1/10 -

1/100)

Entrez Gene: 5970/19697

O04206/O04207 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

WB description:

Western blot analysis of NF-kappaB p65 (pS468) expression in HeLa TNFa-treated (A), NIH3T3 LPS-treated (B), rat brain (C) whole cell lysates.





IHC description:

Immunohistochemical analysis of NF-kappaB p65 (pS468) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.