

Manufaaturar/Su	nnlior	
Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>		
Product name:	Rabbit anti-NF-kappaB p65 Antibody	
Catalog:	DL91000A	
Synonyms:	NFKB3; Transcription factor p65; Nuclear factor	
	NF-kappa-B p65 subunit; Nuclear factor of kappa	
	light polypeptide gene enhancer in B-cells 3	
Immunogen:	KLH-conjugated synthetic peptide encompassing a	
	sequence within the center region of human	
	NF-kappaB p65. The exact sequence is proprietary.	
Form: Liquid	Concentration: 1mg/mL	
r or internet		
Size: 100 ul/50 ul	Host: Rabbit	
Size. 100 ul/30 ul	Host. Rabolt	
Reactivity:	Application: WB, IHC, IP	
•	•••	
Human,Mouse,Rat,Bovine,Monke		



y,Pig


Clonality: Polyc	lonal Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)
Entrez Gene:	5970/19697
SwissProt:	Q04206/Q04207
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87%

Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of NF-kappaB p65 expression in HepG2 (A), MCF7 (B), Raw264.7 (C), rat liver (D) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of NF-kappaB p65 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

