

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-SRSF3 Antibody			
Catalog:	DL91058A			
Synonyms:	SFRS3; SRP20; Serine/arginine-rich splicing factor			
	3; Pre-mRNA	-splicing facto	or SRP20; Splicing	
	factor, arginin	e/serine-rich 3	3	
T				
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the C-term region of human SRSF3. The exact sequence is proprietary.			
	SKSF3. The e	xact sequence	is proprietary.	
Form: Liquid		Concentra	ation: 1mg/mL	
I of my Erquita		Concentra		
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human, Mouse		Applicatio	Application: WB, IHC, IF/IC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	6428/20383
SwissProt:	P84103/P84104
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of SRSF3 expression in HepG2 (A), HeLa (B), BJAB (C) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of SRSF3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of SRSF3 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

