

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-SKP2 Antibody			
Catalog:	DL91069A			
Synonyms:	FBXL1; S-phase	e kinase-ass	sociated protein 2;	
	Cyclin-A/CDK2	2-associated	protein p45; F-box	
	protein Skp2; F-	-box/LRR-r	epeat protein 1; p45skp2	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the C-term region of human SKP2.			
	The exact sequence is proprietary.			
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bbit	
Reactivity: Human, Bovine, Pig		Applicati	on: WB, IHC, IF/IC	




Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	6502
SwissProt:	Q13309
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of SKP2 expression in A673 (A), HeLa (B), K562 (C) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of SKP2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of SKP2 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

