

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-STAT1 (pS727) Antibody

Catalog: DL91109A

Synonyms: Signal transducer and activator of transcription

1-alpha/beta; Transcription factor ISGF-3

components p91/p84

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the C-term region of human STAT1. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Application: WB, IHC, IF/IC, IP

Human, Mouse, Rat, Bovine, Chicke



n,Dog,Monkey,Pig,Sheep

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500), IP (1/10 - 1/100)

Entrez Gene: 6774/20848/25125

P40763/P42227/P52631 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

WB description:

Western blot analysis of STAT3 expression in A549 (A), LO2 (B), SP2/0 (C), NIH3T3 (D), PC12 (E), H9C2 (F) whole cell lysates.






IHC description:

Immunohistochemical analysis of STAT3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of STAT3 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody





IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.