

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Rabbit anti-TSG6 Antibody		
Catalog:	DL91176A		
C			
Synonyms:	TSG6; Tumor necrosis factor-inducible gene 6 protein; Hyaluronate-binding protein; TNF-stimulated gene 6 protein; TSG-6; Tumor		
	necrosis factor alpha-induced protein 6; TNF		
	alpha-induced protein 6		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
8	sequence within the center region of human TSG6.		
	The exact sequence is proprietary.		
	The church bequence is proprietally.		
Form: Liquid	Concentration: 1mg/mL		
Lorino Diquid	Concentration. Img IIL		
Size: 100 ul/50 ul	Host: Rabbit		
5120. 100 ul/ 50 ul	rost. Rubbl		




Reactivity: Human,Bovine,Rabbit		Application: WB, IHC, IP
Clonality: Polyclonal		Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)
Entrez Gene:	7130	
SwissProt:	P98066	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.	

WB description:

Western blot analysis of TSG6 expression in MCF7 (A), HCT15 (B), HeLa (C) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of TSG6 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

