

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-UDG Antibody			
Catalog:	DL91212A			
Synonyms:		UNG15; Ura	cil-DNA glycosylase;	
	UDG			
T		1 (1 (
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the center region of human UDG. The exact sequence is proprietary.			
	The exact sequ	ience is propi	netary.	
Form: Liquid		Concentr	ation: 1mg/mL	
1 01 111 21 1 11 11				
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity:		Applicati	on: WB, IHC	
Human,Mouse,Rat,Bovine,Pig				



Clonality: Polyclonal


Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)

Entrez Gene:	7374/22256
SwissProt:	P13051/P97931
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of UDG expression in HeLa (A), Jurkat (B) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of UDG staining in human muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.