

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Vimentin Antibody			
Catalog:	DL91228A			
Synonyms:	Vimentin			
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Vimentin. The exact sequence is proprietary.			
Form: Liquid		Concentra	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human,Mouse,Rat n,Monkey,Pig	Bovine,Chicke,	Application	on: WB, IHC, IF/IC	



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	7431/22352/81818
SwissProt:	P08670/P20152/P31000
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Vimentin expression in MCF7 (A), MDAMB435 (B), KNRK (C), HeLa (D), Raw264.7 (E), PC12 (F) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of Vimentin staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of Vimentin staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibo

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

