

| Manufacturer/Supplier: | | | | |
|---|---|---------------|------------------------|--|
| WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD | | | | |
| Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | | | |
| Wuxi Jiangsu Province, China | | | | |
| TELE: 86-510-82732223 FAX: 86-510-82720101-8014 | | | | |
| Web : <u>www.dldevelop.com</u> | | | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | | | |
| | | | | |
| Product name: | Rabbit anti-PEX11B Antibody | | | |
| Catalog: | DL91323A | | | |
| | | | | |
| Synonyms: | Peroxisomal membrane protein 11B; Peroxin-11B; | | | |
| | Peroxisomal bi | iogenesis fac | tor 11B; Protein PEX11 | |
| | homolog beta; | PEX11-beta | | |
| | | | | |
| Immunogen: | KLH-conjugated synthetic peptide encompassing a | | | |
| | sequence within the center region of human | | | |
| | PEX11B. The exact sequence is proprietary. | | | |
| | | | | |
| Form: Liquid | | Concentra | ation: 1mg/mL | |
| - | | | - | |
| Size: 100 ul/50 ul | | Host: Rab | bit | |
| | | | | |
| Reactivity: | | Application | on: WB, IHC, IF/IC, IP | |
| Human, Mouse, Rat, Bovine | | | | |
| | | | | |
| | | | | |



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 -1/500), IP (1/10 - 1/100)


| Entrez Gene: | 8799/18632 |
|---------------|---|
| SwissProt: | O96011/Q9Z210 |
| Purification: | The antibody was purified by immunogen affinity chromatography. |
| Buffer: | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |

WB description:

Western blot analysis of PEX11B expression in HT29 (A), THP1 (B), Jurkat (C) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of PEX11B staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of PEX11B staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

DEVELOP.



IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

