

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-CDK5R1 p35 Antibody		
Catalog:	DL91333A		
Synonyms:	CDK5R; NCK5A; Cyclin-dependent kinase 5		
	activator 1; CDK5 activator 1; Cyclin-dependent kinase 5 regulatory subunit 1; TPKII regulatory		
	subunit		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
	sequence within the N-term region of human CDK5R1 p35. The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
CI 100 1/50 1			
Size: 100 ul/50 ul	Host: Rabbit		
Desetivity	Applications WD IE/IC		
Reactivity:	Application: WB, IF/IC		

DEVELOP.


Human, Mouse, Rat, Bovine, Zebraf ish

Clonality: Polyclo	nal	Dilution: WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)
Entrez Gene:	8851/12569/116671	
SwissProt:	Q15078/P61809/P61810	
Purification:	The antibody wa chromatography	as purified by immunogen affinity
Buffer:	1	Potassium phosphate, 0.87% c, pH 7.3, 30% glycerol, and 0.01%

WB description:


Western blot analysis of CDK5R1 p35 expression in rat brain (A), HeLa (B), A431 (C) whole cell lysates.

DEVELOP



IF/ICC description:

Immunofluorescent analysis of CDK5R1 p35 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary anti



Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.

