

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-JAB1 Antibody			
Catalog:	DL91523A			
G	CONT LADI	COD0 : 1	1 1 4	
Synonyms:		C C	osome complex subunit	
			unit 5; Jun activation	
	domain-bindin	ig protein 1		
Immunogen:	KLH-conjugat	ted synthetic i	nentide encompassing a	
minunogen.	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human JAB1.			
	The exact sequence is proprietary.			
	The exact sequ	ience is propi	ictury.	
Form: Liquid		Concentra	ation: 1mg/mL	
1			0	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity:		Applicatio	on: WB, IHC	
Human,Mouse,Rat,Monkey,Pig,Z				




ebrafish


Clonality: Polyclo	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)		
Entrez Gene:	10987/26754		
SwissProt:	Q92905/O35864		
Purification:	The antibody was purified by immunogen affinity chromatography.		
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.019 sodium azide.		

WB description:

Western blot analysis of JAB1 expression in Caki1 (A), mouse embryo (B), rat liver (C) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of JAB1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.