

| Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District | | |
|--------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------|--|
| Wuxi Jiangsu Province, China | | |
| TELE: 86-510-82 | 732223 FAX: 86-510-82720101-8014 | |
| Web : <u>www.dldevelop.com</u> | | |
| Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u> | | |
| Product name: Catalog: | Rabbit anti-RPL36 Antibody DL91602A | |
| Synonyms: | 60S ribosomal protein L36 | |
| Immunogen: | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human RPL36. The exact sequence is proprietary. | |
| Form: Liquid | Concentration: 1mg/mL | |
| Size: 100 ul/50 ul | Host: Rabbit | |
| | | |

Reactivity: Human,Mouse,Rat,Bovine,Chicke Application: WB, IHC, IF/IC



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


| Entrez Gene: | 25873/58927 |
|---------------|-----------------------------------------------------------------------------------------------------------------|
| SwissProt: | Q9Y3U8/P47964/P39032 |
| Purification: | The antibody was purified by immunogen affinity chromatography. |
| Buffer: | Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide. |

WB description:

Western blot analysis of RPL36 expression in MDAMB435 (A), HEK293 (B), Raw264.7 (C), mouse brain (D), rat brain (E) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of RPL36 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of RPL36 staining in MDAMB435 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

