

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-A4GNT Antibody			
Catalog:	DL91660A			
C				
Synonyms:	Alpha-1,4-N-acetylglucosaminyltransferase;			
	Alpha4GnT	58	,	
	i iipiiu toiri			
Immunogen:	KLH-conjugated	d synthetic	peptide encompassing a	
8	sequence within the center region of human			
	A4GNT. The exact sequence is proprietary.			
		det sequent	te is proprietary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Form. Elquid		Concenti	ation. mig/mL	
Size: 100 ul/50 ul		Host: Rab	hit	
Size: 100 ul/30 ul		HUSt. Kat	on	
Reactivity: Human, Mouse, Rat		Applicati	on: WB, IHC	
Reactivity. Human, wouse, Kat		присан	M. WD , MC	
Clonality: Polyclonal		Dilution	WB (1/500 - 1/1000),	
Cionanty: Polyclo	llai	Dilution:	(1/300 - 1/1000),	




IHC (1/100 - 1/200)

Entrez Gene:	51146
SwissProt:	Q9UNA3
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of A4GNT expression in HEK293T (A), mouse testis (B), rat kidney (C) whole cell lysates.





Immunohistochemical analysis of A4GNT staining in human pancreas formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

