

Manufacturer/Supplier: WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82	2732223	FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-ARHGAP18 Antibody			
Catalog:	DL91834A			
cg.	22,100			
Synonyms:	Pha CTDaga activating protain 18, MagCAD			
Synonyms.	Rho GTPase-activating protein 18; MacGAP; Rho-type GTPase-activating protein 18			
	Rho-type GTPa	se-activatin	g protein 18	
-				
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the C-term region of human			
	ARHGAP18. Th	he exact sec	juence is proprietary.	
Form: Liquid		Concentration: 1mg/mL		
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Monkey		Applicati	on: WB, IHC, IP	
<b>Clonality:</b> Polyclonal		Dilution	WB (1/500 - 1/1000),	



IHC (1/100 - 1/200), IP (1/10 - 1/100)


Entrez Gene:	93663
SwissProt:	Q8N392
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of ARHGAP18 expression in MDAMB231 (A), U2OS (B) whole cell lysates.



# **DEVELOP**



### **IHC description:**

Immunohistochemical analysis of ARHGAP18 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IP description:**

### **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.

