

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: info@dldevelop.com.cn service@dldevelop.com.cn

Product name: Rabbit anti-NCOA7 Antibody

Catalog: DL91863A

ERAP140; ESNA1; Nuclear receptor coactivator 7; **Synonyms:**

140 kDa estrogen receptor-associated protein;

Estrogen nuclear receptor coactivator 1

Immunogen: KLH-conjugated synthetic peptide encompassing a

> sequence within the center region of human NCOA7. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 u1/50 u1 Host: Rabbit

Reactivity: Human, Mouse, Rat **Application:** WB, IHC, IP



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IP (1/10 -

1/100)

Entrez Gene: 135112/211329

O8NI08/O6DFV7 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of NCOA7 expression in HEK293T (A), mouse kidney (B), rat kidney (C) whole cell lysates.





IHC description:

Immunohistochemical analysis of NCOA7 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.