

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Province, China	
TELE: 86-510-82	FAX: 86-510-82720101-8014
Web : <u>www.dldevelop.com</u>	
Email: info@dldevelop.com.cn service@dldevelop.com.cn	
Product name:	Rabbit anti-Caspase 9 Antibody
Catalog:	DL92157A
Synonyms:	MCH6; Caspase-9; CASP-9; Apoptotic protease
	Mch-6; Apoptotic protease-activating factor 3;
	APAF-3; ICE-like apoptotic protease 6; ICE-LAP6
Immunogen:	KLH-conjugated synthetic peptide encompassing a
	sequence within the center region of human Caspase
	9. The exact sequence is proprietary.
Former Lincid	
Form: Liquid	Concentration: 1mg/mL
<b>Size:</b> 100 ul/50 ul	Host: Rabbit
Size: 100 ui 20 ui	HOST INCOM
Reactivity: Human	n Application: WB, IHC, IF/IC
v	A.A.



**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene: 842


SwissProt: P55211

**Purification:** The antibody was purified by immunogen affinity chromatography.


**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB** description:

Western blot analysis of Caspase 9 expression in HeLa (A), Jurkat (B) whole cell lysates.



# **DEVELOP**




#### **IHC description:**

Immunohistochemical analysis of Caspase 9 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of Caspase 9 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib

## **DEVELOP**.



### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

