

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: info@dldevelop.com.cn service@dldevelop.com.cn			
Product name:	Rabbit anti-CDK8 Antibody		
Catalog:	DL92171A		
_			
Synonyms:	Cyclin-dependent kinase 8; Cell division protein		
	kinase 8; Mediator complex subunit CDK8;		
	Mediator of RNA polymerase II transcription		
	subunit CDK8; Protein kinase K35		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
U	sequence within the N-term region of human CDK8.		
	The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
1			
Size: 100 ul/50 ul	Host: Rabbit		
<b>Reactivity:</b>	<b>Application:</b> WB, IHC		

# **DEVELOP**

## Human, Mouse, Rat, Zebrafish


Clonality: Polyclo	nal	<b>Dilution:</b> WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Entrez Gene:	1024/264064	
SwissProt:	P49336/Q8R3L8	
Purification:	The antibody wa chromatography.	s purified by immunogen affinity

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of CDK8 expression in HepG2 (A), A549 (B), Raw264.7 (C) whole cell lysates.

# **DEVELOP**



### **IHC description:**

Immunohistochemical analysis of CDK8 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.