

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

Product name: Rabbit anti-MKK2 (pT394) Antibody

Catalog: DL92301A

Synonyms: MEK2; MKK2; PRKMK2; Dual specificity

mitogen-activated protein kinase kinase 2; MAP kinase kinase 2; MAPKK 2; ERK activator kinase

2; MAPK/ERK kinase 2; MEK 2

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the C-term region of human MKK2

(pT394). The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

Reactivity: Human, Monkey Application: WB, IHC, IP



Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IP (1/10 -

1/100)

Entrez Gene: 5605

P36507 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

WB description:

Western blot analysis of MKK2 (pT394) expression in HeLa UV-treated (A), HepG2 UV-treated (B) whole cell lysates.





IHC description:

Immunohistochemical analysis of MKK2 (pT394) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.