

Manufacturer/Supplier:WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTDAddress: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi DistrictWuxi Jiangsu Province, ChinaTELE: 86-510-82732223FAX: 86-510-82720101-8014Web : www.dldevelop.comEmail: info@dldevelop.com.cnservice@dldevelop.com.cn			
Product name:	Rabbit anti-SHPTP2 Antibody		
Catalog:	DL92306A		
Synonyms: Immunogen:	PTP2C; SHPTP2; Tyrosine-protein phosphatase non-receptor type 11; Protein-tyrosine phosphatase 1D; PTP-1D; Protein-tyrosine phosphatase 2C; PTP-2C; SH-PTP2; SHP-2; Shp2; SH-PTP3 KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human SHPTP2. The exact sequence is proprietary.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		
Reactivity:	Application: WB, IHC		

DEVELOP.


Human,Mouse,Rat,Bovine,Chicke n,Dog,Pig,Sheep,Rabbit

Clonality: Polyclo	nal	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Entrez Gene:	5781/19247/25622	
SwissProt:	Q06124/P35235/P41499	
Purification:	The antibody wa chromatography	as purified by immunogen affinity
Buffer:	1	Potassium phosphate, 0.87% , pH 7.3, 30% glycerol, and 0.01%

WB description:

Western blot analysis of SHPTP2 expression in HeLa (A), Raw264.7 (B) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of SHPTP2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.