

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-Tyro	osine Hydro	oxylase (pS71) Antibody	
Catalog:	DL92337A			
Synonyms:	TYH; Tyrosine 3-monooxygenase; Tyrosine			
	3-hydroxylase;	ГН		
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
U	sequence within the N-term region of human			
	Tyrosine Hydroxylase (pS71). The exact sequence			
	is proprietary.	, 4	, 1	
Form: Liquid		Concentra	ation: 1mg/mL	
1			8	
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity:		Applicati	on: WB, IHC	
Human,Mouse,Rat,Chicken,Dog				



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)


Entrez Gene:	7054/21823/25085
SwissProt:	P07101/P24529/P04177
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of Tyrosine Hydroxylase (pS71) expression in HepG2 UV-treated (A), mouse brain (B), PC12 (C) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of Tyrosine Hydroxylase (pS71) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

