

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

**Product name:** Rabbit anti-Histone Deacetylase 8 (pS39) Antibody

Catalog: DL92420A

**Synonyms:** HDACL1; Histone deacetylase 8; HD8

**Immunogen:** KLH-conjugated synthetic peptide encompassing a

sequence within the N-term region of human

Histone Deacetylase 8 (pS39). The exact sequence

is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

**Reactivity:** Application: WB, IHC, IF/IC

Human, Mouse, Rat, Monkey



**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

Entrez Gene: 55869/70315/100911968; 363481

O9BY41/O8VH37/B1WC68 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

## **WB** description:

Western blot analysis of Histone Deacetylase 8 (pS39) expression in A549 (A), NIH3T3 (B), rat kidney (C) whole cell lysates.





## **IHC** description:

Immunohistochemical analysis of Histone Deacetylase 8 (pS39) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IF/ICC description:**

Immunofluorescent analysis of Histone Deacetylase 8 (pS39) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed wit





**Storage:** Store at -20°C. Avoid repeated freeze / thaw cycles.