

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-HECW2 Antibody			
Catalog:	DL92425A			
Synonyms:	KIAA1301; NEDL2; E3 ubiquitin-protein ligase			
	HECW2; HEC	T, C2 and W	W domain-containing	
	protein 2; NED	DD4-like E3	ubiquitin-protein ligase 2	
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	-		region of human	
	HECW2. The	exact sequen	ce is proprietary.	
T		C		
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rab	shit	
SIZC. 100 ul/30 ul		nost. Rat	.010	
Reactivity: Human, Mouse		Applicati	Application: WB, IF/IC	



Clonality: Polyclonal


Dilution: WB (1/500 - 1/1000), IF/IC (1/100 - 1/500)

Entrez Gene:	57520/329152
SwissProt:	Q9P2P5/Q6I6G8
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:


Western blot analysis of HECW2 expression in A549 (A), NIH3T3 (B) whole cell lysates.

DEVELOP



IF/ICC description:

Immunofluorescent analysis of HECW2 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Storage:



Store at -20°C. Avoid repeated freeze / thaw cycles.

