

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223		FAX:	86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-MRPL11 Antibody			
Catalog:	DL92431A			
g.				
Synonyms:	39S ribosomal protein L11, mitochondrial; L11mt;			
Synonyms.	MRP-L11		initoenonariai, E11int,	
	WINF-LIII			
Immunogon	VI U conjugato	downthatia	peptide encompassing a	
Immunogen:		•		
	-		region of human	
	MRPL11. The e	xact sequer	nce is proprietary.	
		_		
Form: Liquid		Concentr	ation: 1mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
Reactivity: Human, Monkey		Applicati	on: WB, IHC, IF/IC	
Clonality: Polyclonal		Dilution:	WB (1/500 - 1/1000),	




IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene:	65003
SwissProt:	Q9Y3B7
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of MRPL11 expression in A431 (A), HepG2 (B) whole cell lysates.

DEVELOP




IHC description:

Immunohistochemical analysis of MRPL11 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of MRPL11 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

