

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510			86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-NT5C1B Antibody			
Catalog:	DL92467A			
Synonyms:	AIRP; Cytosolic 5'-nucleotidase 1B; cN1B;			
	Autoimmune in	fertility-rela	ted protein; Cytosolic	
	5'-nucleotidase	IB; cN-IB		
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
sequence within the center region of human			0	
	NT5C1B. The e	xact sequen	ce is proprietary.	
Form: Liquid		Concentr	ation: 1mg/mL	
Form. Liquid		Concentra		
Size: 100 ul/50 ul		Host: Rab	bit	
Reactivity: Human		Application: WB, IHC, IF/IC		



Clonality: Polyclonal


Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene:


SwissProt:	Q96P26
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of NT5C1B expression in EAhy926 (A) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of NT5C1B staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of NT5C1B staining in EAhy926 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

