

**Manufacturer/Supplier:**

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

**Address:** A1-203 Mingpin CityII, No.8 Xihudong Road, Liangxi District  
Wuxi Jiangsu Province, China

TELE: 86-510-82732223

FAX: 86-510-82720101-8014

**Web :** [www.dldevelop.com](http://www.dldevelop.com)

Email: [info@dldevelop.com.cn](mailto:info@dldevelop.com.cn)    [service@dldevelop.com.cn](mailto:service@dldevelop.com.cn)

**Product name:** Rabbit anti-AKT Antibody

**Catalog:** DL92759A

**Synonyms:** AKT1; PKB; RAC; RAC-alpha  
serine/threonine-protein kinase; Protein kinase B;  
PKB; Protein kinase B alpha; PKB alpha;  
Proto-oncogene c-Akt; RAC-PK-alpha; AKT2;  
RAC-beta serine/threonine-protein kinase; Protein  
kinase Akt-2; Protein kinase B beta; PKB beta;

**Immunogen:** KLH-conjugated synthetic peptide encompassing a  
sequence within the center region of human AKT.  
The exact sequence is proprietary.

**Form:** Liquid

**Concentration:** 1mg/mL

**Size:** 100 ul/50 ul

**Host:** Rabbit

**Reactivity:**

Human, Mouse, Rat, Bovine, Sheep

**Application:** WB, IHC, IF/IC**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000),  
IHC (1/100 - 1/200), IF/IC (1/100 -  
1/500)

**Entrez Gene:**

11651; 11652; 23797/24185; 25233; 29414

**SwissProt:**

P31749; P31751; Q9Y243/P31750; Q60823;  
Q9WUA6/P47196; P47197; Q63484

**Purification:**


The antibody was purified by immunogen affinity chromatography.

**Buffer:**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

**WB description:**

Western blot analysis of AKT expression in MCF7 (A), NIH3T3 (B) whole cell lysates.




### IHC description:

Immunohistochemical analysis of AKT staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### IF/ICC description:

Immunofluorescent analysis of AKT staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in



**Storage:**

Store at -20°C. Avoid repeated freeze / thaw cycles.