


Manufacturer/Supplier:WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTDAddress: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi DistrictWuxi Jiangsu Province, ChinaTELE: 86-510-82732223FAX: 86-510-82720101-8014Web : www.dldevelop.comEmail: info@dldevelop.com.cnservice@dldevelop.com.cn		
Product name: Catalog:	Rabbit anti-BAD (pS112) Antibody DL92792A	
Synonyms:	BBC6; BCL2L8; Bcl2 antagonist of cell death; BAD; Bcl-2-binding component 6; Bcl-2-like protein 8; Bcl2-L-8; Bcl-XL/Bcl-2-associated death promoter	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human BAD (pS112). The exact sequence is proprietary.	
Form: Liquid	Concentration: 1mg/mL	
Size: 100 ul/50 ul	Host: Rabbit	
Reactivity: Huma	n Application: WB, IHC	




Clonality: Polyclon	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200)
Entrez Gene:	572
SwissProt:	Q92934
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of BAD (pS112) expression in HeLa Forskolin-treated (A), MCF7 (B) whole cell lysates.



DEVELOP



IHC description:

Immunohistochemical analysis of BAD (pS112) staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.