

## Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

**Product name:** Rabbit anti-BIM (pS69) Antibody

Catalog: DL92802A

**Synonyms:** BIM; Bcl-2-like protein 11; Bcl2-L-11;

Bcl2-interacting mediator of cell death

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human BIM

(pS69). The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit

**Reactivity:** Application: WB, IHC, IF/IC

Human, Mouse, Rat, Bovine, Sheep



**Clonality:** Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200), IF/IC (1/100 -

1/500)

Entrez Gene: 10018/12125/64547

O43521/O54918/O88498 SwissProt:

**Purification:** The antibody was purified by immunogen affinity

chromatography.

**Buffer:** Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide.

**WB** description:

Western blot analysis of BIM (pS69) expression in HEK293T (A), Raw264.7 (B) whole cell lysates.






## **IHC** description:

Immunohistochemical analysis of BIM (pS69) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

## **IF/ICC description:**

Immunofluorescent analysis of BIM (pS69) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary anti





**Storage:** Store at -20°C. Avoid repeated freeze / thaw cycles.