

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>				
Product name:	Rabbit anti-CaMK1 alpha (pT177) Antibody			
Catalog:	DL92811A			
Synonyms:	Calcium/calmodulin-dependent protein kinase type			
	1; CaM kinase I	; CaM-KI; Cal	M kinase I alpha;	
	CaMKI-alpha			
Immunogen:	KLH-conjugated synthetic peptide encompassing a			
	sequence within the center region of human CaMK1			
	alpha (pT177).	The exact sequ	ence is proprietary.	
Forme Liquid		Concentratio	one 1m a/mI	
Form: Liquid		Concentratio	on: mg/mL	
Size: 100 ul/50 ul		Host: Rabbit		
SHEET TOO MI DO MI				
Reactivity: Human, Mouse, Rat		Application: WB, IHC, IF/IC		



**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	8536/52163/171503
SwissProt:	Q14012/Q91YS8/Q63450
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of CaMK1 alpha (pT177) expression in Jurkat Insulin-treated (A) whole cell lysates.



# **DEVELOP**



#### **IHC description:**


Immunohistochemical analysis of CaMK1 alpha (pT177) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **IF/ICC description:**

Immunofluorescent analysis of CaMK1 alpha (pT177) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the p



## **DEVELOP**.



### Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

