

Manufacturer/Supplier:	
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD	
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District	
Wuxi Jiangsu Province, China	
TELE: 86-510-82732223 FAX: 86-510-82720101-8014	
Web : <u>www.dldevelop.com</u>	
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>	
Product name:	Rabbit anti-COX4-1 Antibody
Catalog:	DL92866A
Synonyms:	COX4; Cytochrome c oxidase subunit 4 isoform 1,
	mitochondrial; Cytochrome c oxidase polypeptide
	IV; Cytochrome c oxidase subunit IV isoform 1;
	COX IV-1
Immunogen:	KLH-conjugated synthetic peptide encompassing a
C	sequence within the N-term region of human
	COX4-1. The exact sequence is proprietary.
Form: Liquid	Concentration: 1mg/mL
1	5
Size: 100 ul/50 ul	Host: Rabbit
Reactivity: Human	n Application: WB, IHC, IF/IC



Clonality: Polyclonal

Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)

Entrez Gene: 1327


SwissProt: P13073

Purification: The antibody was purified by immunogen affinity chromatography.


Buffer: Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of COX4-1 expression in HepG2 (A), Jurkat (B) whole cell lysates.



DEVELOP




IHC description:

Immunohistochemical analysis of COX4-1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of COX4-1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody

DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

