

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>		
Rabbit anti-DLX3 Antibody		
DL92900A		
Homeobox protein DLX-3		
KLH-conjugated synthetic peptide encompassing a		
sequence within the center region of human DLX3.		
The exact sequence is proprietary.		
Concentration: 1mg/mL		

Size: 100 ul/50 ul

Host: Rabbit

Reactivity:Application: WB, IHC, IF/ICHuman,Mouse,Rat,Bovine,Dog,Pig,Sheep




**Clonality:** Polyclonal

**Dilution:** WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/IC (1/100 - 1/500)


Entrez Gene:	1747/13393
SwissProt:	O60479/Q64205
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### **WB description:**

Western blot analysis of DLX3 expression in JAR (A), rat skin (B) whole cell lysates.



# **DEVELOP**



#### **IHC description:**

Immunohistochemical analysis of DLX3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### **IF/ICC description:**

Immunofluorescent analysis of DLX3 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



# **DEVELOP**.



## Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

