

Manufacturer/Supplier:				
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD				
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District				
Wuxi Jiangsu Province, China				
TELE: 86-510-82732223 FAX: 86-510-82720101-8014				
Web : <u>www.dldevelop.com</u>				
Email: info@dldevelop.com.cn service@dldevelop.com.cn				
Product name:	Rabbit anti-JUN	B Antibody	Į	
Catalog:	DL93043A			
cuturog.	DEFECTION			
Sumonymer	Transcription factor jun-B			
Synonyms:	Transcription fa	CIOI JUII-D		
Immunagan	KI II conjugated synthetic nontide encomposing a			
mmunogen.				
	1		e	
	The exact seque	nce is propi	netary.	
		<b>C</b>		
Form: Liquid		Concentr	ation: Img/mL	
Size: 100 ul/50 ul		Host: Rab	obit	
Reactivity:		Applicati	on: WB, IHC, IF/IC, IP	
Human,Mouse,Rat,Bovine,Sheep				
<b>Clonality:</b> Polyclonal		<b>Dilution:</b> WB (1/500 - 1/1000),		
sequence within The exact sequen Form: Liquid Size: 100 ul/50 ul Reactivity: Human,Mouse,Rat,Bovine,Sheep		d synthetic peptide encompassing a the C-term region of human JUNB. nce is proprietary. Concentration: 1mg/mL Host: Rabbit Application: WB, IHC, IF/IC, IP Dilution: WB (1/500 - 1/1000),		




IHC (1/100 - 1/200), IF/IC (1/100 - 1/500), IP (1/10 - 1/100)

Entrez Gene:	4791/18034
SwissProt:	Q00653/Q9WTK5
Purification:	The antibody was purified by immunogen affinity chromatography.
Buffer:	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

### WB description:

Western blot analysis of NF-kappaB p100 (pS865) expression in HeLa (A), MCF7 (B) whole cell lysates.

# DEVELOP




#### **IHC description:**

Immunohistochemical analysis of NF-kappaB p100 (pS865) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

### **IF/ICC description:**

Immunofluorescent analysis of NF-kappaB p100 (pS865) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the

# **DEVELOP**.



**IP description:** 

**Storage:** 

Store at -20°C. Avoid repeated freeze / thaw cycles.

