

Manufacturer/Supplier:

WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD

Address: A1-203 Mingpin CityII, No.8 Xihudong Road. Liangxi District

Wuxi Jiangsu Province, China

TELE: 86-510-82732223 FAX: 86-510-82720101-8014

Web: www.dldevelop.com

Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>

Product name: Rabbit anti-TRAP230 Antibody

Catalog: DL93084A

Synonyms: ARC240; CAGH45; HOPA; KIAA0192; TNRC11;

TRAP230; Mediator of RNA polymerase II transcription subunit 12; Activator-recruited

cofactor 240 kDa component; ARC240; CAG repeat

protein 45; Mediator complex subunit 12; OPA-containing protein; Thyroid hormone re

Immunogen: KLH-conjugated synthetic peptide encompassing a

sequence within the center region of human TRAP230. The exact sequence is proprietary.

Form: Liquid Concentration: 1mg/mL

Size: 100 ul/50 ul Host: Rabbit



Reactivity: Human, Mouse, Rat **Application:** WB, IHC

Clonality: Polyclonal **Dilution:** WB (1/500 - 1/1000),

IHC (1/100 - 1/200)

Entrez Gene: 9968/59024

O93074/A2AGH6 SwissProt:

Purification: The antibody was purified by immunogen affinity

chromatography.

Buffer: Liquid in 0.42% Potassium phosphate, 0.87%


Sodium chloride, pH 7.3, 30% glycerol, and 0.01%

sodium azide

WB description:

Western blot analysis of TRAP230 expression in HEK293T (A), NS-1 (B), PC12 (C) whole cell lysates.





IHC description:

Immunohistochemical analysis of TRAP230 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.