

Manufacturer/Supplier:		
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD		
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District		
Wuxi Jiangsu Province, China		
TELE: 86-510-82732223 FAX: 86-510-82720101-8014		
Web : <u>www.dldevelop.com</u>		
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>		
Product name:	Rabbit anti-hnRNP G Antibody	
Catalog:	DL93195A	
Synonyms:	HNRPG; RBMXP1; RNA-binding motif protein, X	
	chromosome; Glycoprotein p43; Heterogeneous	
	nuclear ribonucleoprotein G; hnRNP G	
Immunogen:	KLH-conjugated synthetic peptide encompassing a	
	sequence within the N-term region of human	
	hnRNP G. The exact sequence is proprietary.	
Form: Liquid	Concentration: 1mg/mL	
C* 100 1/50 1		
Size: 100 ul/50 ul	Host: Rabbit	
Reactivity:	Application: WB, IHC, IF/IC	
Human,Mouse,Rat,Bovine,Pig,Ze		
11umun,1110u00,11ut,1001110,115,20		



brafish

Clonality: Polyclonal	Dilution: WB (1/500 - 1/1000),
	IHC (1/100 - 1/200), IF/IC (1/100 -
	1/500)

Entrez Gene: 27316/19655/302855


SwissProt: P38159/Q9WV02/Q4V898

- **Purification:** The antibody was purified by immunogen affinity chromatography.
- **Buffer:** Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

WB description:

Western blot analysis of hnRNP G expression in Jurkat (A), RT4 (B) whole cell lysates.

DEVELOP



IHC description:


Immunohistochemical analysis of hnRNP G staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IF/ICC description:

Immunofluorescent analysis of hnRNP G staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antib



DEVELOP.



Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

