

Manufacturer/Supplier:			
WUXI DONGLIN SCI&TECH DEVELOPMENT CO., LTD			
Address: A1-203 Mingpin CityII,No.8 Xihudong Road.Liangxi District			
Wuxi Jiangsu Province, China			
TELE: 86-510-82732223 FAX: 86-510-82720101-8014			
Web : <u>www.dldevelop.com</u>			
Email: <u>info@dldevelop.com.cn</u> <u>service@dldevelop.com.cn</u>			
Product name:	Rabbit anti-NF-kappaB p65 (pT254) Antibody		
Catalog:	DL93197A		
C	NEVD2 Township for the second for Northern for the		
Synonyms:	NFKB3; Transcription factor p65; Nuclear factor NF-kappa-B p65 subunit; Nuclear factor of kappa		
	light polypeptide gene enhancer in B-cells 3		
Immunogen:	KLH-conjugated synthetic peptide encompassing a		
g	sequence within the center region of human		
	NF-kappaB p65 (pT254). The exact sequence is		
	proprietary.		
Form: Liquid	Concentration: 1mg/mL		
Size: 100 ul/50 ul	Host: Rabbit		
Reactivity:	Application: WB, IHC, IP		

DEVELOP.


Human,Mouse,Rat,Bovine,Monke y,Pig

Clonality: Polyclo	nal	Dilution: WB (1/500 - 1/1000), IHC (1/100 - 1/200), IP (1/10 - 1/100)
Entrez Gene:	5970/19697	
SwissProt:	Q04206/Q04207	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Buffer:	1	Potassium phosphate, 0.87% , pH 7.3, 30% glycerol, and 0.01%

WB description:

Western blot analysis of NF-kappaB p65 (pT254) expression in HeLa TNFa-treated (A), HL60 TNFa-treated (B) whole cell lysates.

DEVELOP



IHC description:

Immunohistochemical analysis of NF-kappaB p65 (pT254) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

IP description:

Storage:

Store at -20°C. Avoid repeated freeze / thaw cycles.

